arXiv:1404.5820v1 [astro-ph.CO] 23 Apr 2014

LIGHT MILLICHARGED PARTICLES AND LARGE SCALE COSMIC MAGNETIC FIELDS

Alexander D. Dolgov [a]
Novosibirsk State University, Novosibirsk, 630090, Russia
Institute of Theoretical and Experimental Physics, Moscow, 113259, Russia
Dipartimento di Fisica, Università degli Studi di Ferrara, I-44100 Ferrara, Italy

Abstract. After a brief review of different types of scenarios suggested for the solution of the problem of galactic and intergalactic magnetic field generation, the mechanism based on the electric current induction by hypothetical millicharged particles interacting with electrons in cosmic medium is discussed. The proposed model successfully describes observational data. The new light millicharged particles can contribute from a small fraction up to 100% to the cosmological dark matter.

There are strong observational observational evidence for existence of galactic magnetic fields with the field strength B_{gal} of a few μG and the coherence length of several kiloparsecs. There are also some data in favor of weaker intergalactic magnetic fields, $B \sim 10^{-9}$ G, but with much larger coherence scale of hundreds kiloparseks. For a review on observations see refs. [1]. The origin of such large scale galactic and intergalactic magnetic fields remains a cosmological mystery for many years.

There are a lot of proposals for mechanisms of large scale magnetic field generation. They can be roughly divided into two classes: either ones based on conventional physics, or those which invoke different new physics hypotheses. They are reviewed e.g. in papers [2]. Both types of mechanisms suffer from the similar problems. The magnitude of the generated field might be quite high but the coherence length happened to be much shorter than the necessary one, or vice versa: the coherence length is large but the magnitude is tiny. The problems are milder in the case of new physics allowed but they still remain vital. The case of tiny but the galactic scale seed magnetic field to some extend could be cured by the galactic dynamo amplification. However, the seed fields are usually so weak that the galactic dynamo is not sufficient to amplify them up to the observed magnitude. As for intergalactic fields, the dynamo practically is not efficient there, so the existence of intergalactic fields puts strong restrictions on the mechanisms of generation of primordial (seed) magnetic fields.

It is tempting to search for a possibility to generate primordial magnetic fields during inflation, since the problem of large scale in this case is naturally and easily solved by exponential stretching. However, an electromagnetic field cannot be created during inflation because the classical electrodynamics is conformally invariant and such fields are not produced in conformally flat Friedmann-Robertson-Walker metric [3]. However, quantum triangle anomaly

[a] E-mail: dolgov@fe.infn.it

breaks conformal invariance and allows for photon production and thus for magnetic field generation [4]. This is a promising mechanism but more detailed investigation of renormalization in De Sitter space-time is necessary. Another, more exotic possibility, is an assumption of non-minimal coupling of electromagnetic field to gravity through a new interaction, e.g. by term $\xi R A_\mu A^\mu$ [5], where R is the curvature scalar and ξ is a constant. Such an interaction which not only breaks conformal but the gauge invariance of electrodynamics as well. As a result the photon would acquire non-zero mass, due to one-loop graviton-photon diagram, of the order $m_\gamma \sim \xi\Lambda^2/m_{Pl}$, where Λ is an ultraviolet cut-off and $m_{Pl} \approx 10^{19}$ GeV is the Planck mass. An existence of the galactic magnetic fields implies $m_\gamma < 1/\mathrm{kpc}$, which in turn demands $\Lambda < 10\,\mathrm{eV}/\xi$. So for a reasonable ultraviolet cutoff a tiny ξ is necessary which would drastically diminish the effect of this new interaction on generation of primordial magnetic field.

At later cosmological stages very strong magnetic fields could be created at first order phase transitions but the coherence length in this case would be microscopically small and even with cosmological expansion it would remain by far too short. This restriction is avoided if e.g. at electroweak phase transition a condensate of electrically charged W-bosons were formed [6]. In this case the primeval plasma could be spontaneously magnetized inside macroscopically large domains and such magnetic fields may be the seeds for the observed today galactic and intergalactic fields.

Next interesting period for generation of primordial magnetic fields might be the epoch of big bang nucleosynthesis at $t \geq 1$ sec, $T \leq 1$ MeV. The coherence scale in this case is bounded form above $l_c < 100$ pc in terms of the present day units. In the case of big and inhomogeneous primordial lepton asymmetry the magnitude of the the created magnetic field may be sufficiently large, such that after chaotic field line reconnection (analogous to the Brownian motion) the coherence scale might extend to galactic, but not to intergalactic scales [7]. This process proceeds at the expense of a decrease of the field amplitude and galactic dynamo is very much in order.

During the last decade an attention was attracted to generation of cosmic magnetic field during or around the epoch of recombination, $z \sim 1000$, or even closer to the present day. at the period of the large scale structure formation. Magnetic field can be produced by vortex currents, but vorticity perturbations are absent in the primordial density perturbations. However, as it was found in refs. [8], vorticity may be created by the photon diffusion in the second order in the usual scalar, non-vortex, perturbations.

Still in this case and other cases mentioned above a huge, sometimes even unrealistic, dynamo amplification by galactic rotation is necessary. This could help to amplify galactic magnetic fields from originally weak seed fields, but fails for intergalactic magnetic field.

In our recent work [9] we suggested a new mechanism of magnetic field gener-

ation in the universe with already formed first large scale structures, assuming that there exist light millicharged particles. Let us consider first a protogalaxy rotating in the background of the CMB photons. The force exerted by photons on electrons is much larger than that exerted on protons, $F \sim \sigma \sim 1/m^2$, where σ is the cross-section of elastic γe or γp scattering. The difference between e and p accelerations is even bigger, because $a \sim F/m \sim 1/m^3$. So a circular electric current, proportional to the rotational velocity of the protogalaxy, v_{rot}, must be induced. The force acting on electrons is given by the expression: $\vec{F} \sim \vec{v}\sigma_{e\gamma} n_\gamma \omega_\gamma$. This force coherently acts on electrons during the collision time determined by ep-scattering, which is equal to:

$$\tau_{ep} = \frac{m_e^2 \langle v_e^2 \rangle}{4\pi\alpha^2 \langle 1/v_e \rangle n_e L_e} \simeq \frac{m_e^{1/2} T_e^{3/2}}{4\pi\alpha^2 n_e L_e}\,, \tag{1}$$

where the Coulomb logarithm is $L_e \sim 10$ and $\langle v_e^2 \rangle = T_e/m_e$. So we obtain the standard expression for the conductivity: $\kappa = e^2 n_e \tau_{ep}/2m_e \simeq T_e^{3/2}/8\pi\alpha L_e m_e^{1/2}$. The conductivity does not depend on the density of the charge carriers, n_e, unless the latter is so small that the resistance is dominated by neutral particles.

Thus the difference between rotational velocities of e and p is $\Delta v_e = \tau_{ep} F/2m_e$ and the current $j = e n_e \Delta v_e$. Naively estimating B by the Biot-Savart law as $B \sim 4\pi j R$ where R is the galaxy radius, we find that for a typical galaxy with $R \sim 10$ kpc $v_{\rm rot} \sim 100$ km/s: $B \sim \mu$G, very close to the observed value without any dynamo. However, this is incorrect since the time to reach stationary (Bio-Savart) limit is longer than the cosmological time.

To proceed further we need to use the MHD equation modified by presence of an external force:

$$\partial_t \vec{B} = \vec{\nabla}\times\vec{F}/e \,+\, \vec{\nabla}\times(\vec{v}\times\vec{B}) + (4\pi\kappa)^{-1}(\Delta\vec{B} - \partial_t^2\vec{B}). \tag{2}$$

In the limit of high conductivity, the second term in the equation, the advection term, can lead to dynamo amplification of magnetic seed fields once the value of the latter is non-zero.

Assuming $B = 0$ at $t = 0$, we find $\vec{B}(t) = \int_0^t dt\, \vec{\nabla}\times\vec{F}/e$. The largest value of the magnetic seed is generated around the hydrogen recombination at $z_{\rm rec} \sim 10^3$, or $t_{\rm rec} \sim 5 \times 10^5$ yr. Earlier the plasma was strongly coupled and the relative motion of electrons and protons was negligible.

The seed field generated at this epoch with coherence length $\lambda \sim 1$ kpc, corresponding to the present scale of a typical galaxy ~ 1 Mpc, is very weak

$$B_\lambda \sim \Omega_\lambda t_{\rm rec} B_F(t_{\rm rec}) \leq 10^{-20}\, G, \tag{3}$$

where $\Omega_\lambda = |\vec{\nabla}\times\vec{v}|_\lambda \leq 10^3(\delta T/T)^2/\lambda$. The seeds with the coherence length of a few kpc and $B_{\rm seed} > 10^{-15}$ G are needed to fit the observations.

Now we consider the case when instead of CMB photons new DM particles, X collide with protons and electrons and similarly generate an electric current. The current is proportional to the cross-section of Xe-elastic scattering, σ_{Xe}, to n_X/n_e ratio, and to X-particle momentum, $p_X = m_X v_{rot}$. Therefore, to produce stronger than CMB force on electrons, σ_{Xe} should be large. This is possible if X have long range interaction leading to an enhancement of σ_{Xe} at low momentum transfer. So we consider millicharged particles with the mass from a few keV to several MeV.

The bounds on the X-particle charge, $e' = \epsilon e$ are summarized in ref. [9]. If $m_X < m_e$, then from ortho-positronium invisible decays follows $\epsilon < 3.4 \cdot 10^{-5}$. For $m_X = 1$ MeV: $\epsilon < 4.1 \times 10^{-4}$. For $m_X = 100$ MeV: $\epsilon < 5.8 \times 10^{-4}$. We assume that $m_X > 10$ keV to avoid strong limits on e' from the stellar evolution. The BBN bounds can be relaxed if the lepton asymmetry is non-zero [10].

If X-particles were thermally produced, their abundance could be calculated according to the Zeldovich [11] (a decade later called Lee-Weinberg) equation:

$$\Omega_X h^2 \approx 0.023 \; x_f \, g_{*f}^{-1/2} \left(\frac{v\sigma_{\rm ann}}{1 \text{ pb}} \right)^{-1}, \qquad (4)$$

where $x_f \equiv m_X/T_f = 10 + \ln[(g_X \, x_f^{1/2} m_X)/9 g_{*f} \, MeV)]$, g_X is the number of the spin states of X-particle, and g_{*f} is the effective number of particle species in the plasma at $T = T_f$. If $m_X < m_e$, X-particles can annihilate only into photons with $v\sigma(X\bar{X} \to 2\gamma) = \pi\alpha'^2/m_X^2$, where $\alpha' = e'^2/4\pi = \epsilon^2\alpha$. Thus $\Omega_X h^2 \approx 150 \left(10^{-5} m/\epsilon^2 \, \text{keV}\right)^2$. Hence X's would be overproduced if $\epsilon < 3.4 \cdot 10^{-5}$. Additional annihilation into $\bar{\nu}\nu$ or dark photons could help, even if the CMB constraint is fulfilled: $\Omega_X h^2 < 0.007$ [12].

If $m_X > m_e$, then the channel $X\bar{X} \to e^+e^-$ is open and $v\sigma(X\bar{X} \to e^+e^-) = \pi\alpha\alpha'/m_X^2$. Correspondingly: $\Omega_X h^2 = 0.012 \left(10^{-5} m/\epsilon \, \text{MeV}\right)^2$ and e.g. for $m_X = 10$ MeV and $\epsilon = 3 \cdot 10^{-5}$, X-particles can make all DM. Nevertheless Ω_X wlll be taken as free parameter.

The force from X-particles on electrons is $F = \sigma_{eX} \, v_{rel} \, n_X \, m_X \, v_{rot}$, where $v_{rel}\sigma_{eX} = 4\pi\alpha\alpha' L/m_X^2 v_{rel}^3$, $m_X n_X = 10 \, \Omega_X h^2 \kappa(z)(1+z)^3 \, \text{keV/cm}^3$, and $\kappa(z)$ is the dark matter overdensity in galactic halo with respect to the mean cosmological density at redshift z.

Before discussing the generation of B by X-particles let us comment on their role in the structure formation. Prior to recombination the characteristic scattering time of light X-particles is shorter than the universe age, $\tau_{Xe} < t_U$, so they are frozen in $e\gamma$-liquid. After recombination and till reionization they behave as the usual WDM. After reionization τ_{Xe} again becomes shorter than t_U and the rotating ordinary matter in a protogalaxy would transfer a part of angular momentum to X-particles and involve it in its turbulent motion. So at this stage X-particles behave similarly to the usual matter.

For an estimate of the magnitude of B generated by light X-particles we use the obtained above equations but integrate them till reionization, $z = 6$ or $t_u = 1$ Gyr. We take $R = 100$ kpc, $\kappa = 100$, $v_{rot} = 10$ km/sec, and impose the limit $\Omega_x h^2 = 0.007$ to find $B = 10^{-11}\,\mathrm{G}\,\epsilon_5^2/m_{keV}^2$. B can rise by factor 100, becoming 10^{-9} G, when the protogalaxy shrinks from 100 kpc to 10 kpc, by far larger then the minimal necessary strength of the seed.

For heavier X, $m_X > m_e$, a larger charge is allowed, $\epsilon > 10^{-4}$, and X-particles can make all dark matter. After reionization, electron scatterings would not force X-particles into the galaxy rotation and thus the effective integration time can be longer and magnetic fields as large as 10^{-9} G can be generated.

To conclude, we have shown that an existence of millicharged particles with mass in keV - MeV range allows to:
1. Explain the origin of galactic and intergalactic magnetic fields.
2. Introduce DM with time dependent interaction with normal matter.
3. To solve or smooth down the problems of galactic satellites, angular momentum, and cusps in galactic centres inherent to ΛCDM-cosmology.
4. The model can be tested in direct experiment.

Acknowledgment

The author acknowledge the support of the Russian Federation Government Grant No. 11.G34.31.0047.

[1] P.P. Kronberg, Rept. Prog. Phys. **57**, 325 (1994);
J.L. Han, R. Wielebinski, Chinese Journal of Astronomy and Astrophysics, **2**, 293 (2002);
F. Govoni, L. Feretti, Int. J. Mod. Phys. D**13**, 1549 (2004);
R. Beck, AIP Conf. Proc. **1085**, 83 (2009);
A. Neronov, I. Vovk, Science **328**, 523 (2010);
F. Tavecchio *et al.*, MNRAS Lett. **406** L70 (2010);
K. Dolag et al, Astrophys. J. Lett. **727** L4 (2011);
A. M. Taylor, I. Vovk, A. Neronov, Astron. Astrophys. **529** A144 (2011);
K. Takahashi, M. Mori, K. Ichiki, S. Inoue, Astrophys. J. Lett, Volume **744**, L7 (2012);
K. Takahashi, M. Mori, K. Ichiki, S. Inoue, H. Takami, Astrophys. J. Lett, Volume **771**, L42 (2013).

[2] D. Grasso, H.R. Rubinstein, Phys. Repts. **348**, 163 (2001);
A.D. Dolgov, in *From integrable models to gauge theories*, eds. V.G. Gurzadyan *et al.*, pp. 143-154 [hep-ph/0110293];
L.M. Widrow, Rev. Mod. Phys. **74** , 725 (2002);
A. Brandenburg, K. Subramanian, Phys. Repts. **417**, 1 (2005) ;
R.M. Kulsrud, E.G. Zweibel; Rept. Prog. Phys. **71**, 0046091 (2008);
A.D. Dolgov, astro-ph/0306443;

M. Giovannini, Int. J. Mod. Phys. D **13**, 391 (2004);
A. Kandus, K.E. Kunze, C.G. Tsagas, Phys. Repts. **505**, 1 (2011);
L.M. Widrow *et al.*, arXiv:1109.4052.
[3] L. Parker, Phys. Rev. Lett. **21**, 562 (1968).
[4] A.D. Dolgov, Sov. Phys. JETP, bf 54, 223 (1981) [ZETF. **81**, 417 (1981);
A.D. Dolgov, Phys. Rev. D **48**, 2499 (1993).
[5] M.S. Turner, L.M. Widrow, Phys. Rev. D **37**, 2743 (1988).
[6] A.D. Dolgov, A. Lepidi, G. Piccinelli, JCAP **1008**, 031 (2010).
[7] A.D. Dolgov, D. Grasso, Phys. Rev. Lett. **88**, 011301 (2002).
[8] Z. Berezhiani, A.D. Dolgov, Astropart. Phys. **21**, 59 (2004);
S. Matarrese, S. Mollerach, A. Notari, A. Riotto, Phys. Rev. D **71**, 043502 (2005)
[9] Z. Berezhiani, A.D. Dolgov, I.I. Tkachev, Eur.Phys.J. C **73**, 2620 (2013).
[10] Z. Berezhiani, A. Dolgov and I. Tkachev, JCAP **1302**, 010 (2013).
[11] Ya. B. Zeldovich, Adv. Astron. Ap. **3**, 42 (1965).
[12] S.L. Dubovsky, D.S. Gorbunov and G.I. Rubtsov, JETP Lett. **79** (2004);
A.D. Dolgov, S.L. Dubovsky, G.I. Rubtsov, I.I. Tkachev, Phys.Rev. D **88**, 117701 (2013).